

Review Article

Amniotic Membrane in Surgery

M.A. Ganatra

Department of Plastic Surgery, Dow Medical College and Civil Hospital, Karachi.

The first reported use of fetal membrane as skin substitute was by Davis in 1910.¹ In 1913, Salbella presented the first clinical report of successful use of amniotic membrane in the treatment of burns and skin ulcerations.² In 1940, DeRoth reported the use of amniotic membrane in the repair of conjunctival defects.

From 1940 to 1965 a number of clinical trials of successful use of amniotic membrane for use in acute skin injuries appear in the literature.³⁻⁷ However no practical methods of preparation, sterilization and storage were suggested and this fact seems to have limited the use of this modality prior to 1965. In 1965 Dino et al⁸ demonstrated that amniotic membrane from routine deliveries could be sterilized and kept for six weeks at 4°C and safely used on acute second degree burns and on skin donor sites. This encouraging report stimulated great interest amongst clinicians and has resulted in numerous reports in the world literature documenting thousands of patients with successful healing of all kinds of skin lesions.

Amnion is a thin, tough, transparent membrane. It is about 10-15 micrometer thick. It is made up of two membranes, the inner amniotic membrane and the outer chorion. Chorionic side of the membrane is rougher and mucinous. Amnion can easily be separated from chorion leave and placenta as far as the umbilical cord. Once separated, the amnion is found to be smooth and shining and much tougher and more elastic and easier to clean than the thicker chorion, which does not strip from the placenta.¹ The chorion, although thicker, is more easily torn because it is much less elastic.

It is not entirely clear whether the amniotic membrane is primarily nourished by the amniotic fluid or diffusion from the chorion, but the presence of significant ATP and glycogen in the amniotic fluid suggests that the latter is the main source of nourishment of the amniotic membrane.⁹

Immunogenicity of Amniotic Membrane

Amniotic membrane has low or no antigenicity, a fact that might be related apparently to a distinct collagen present in the amniotic membrane. McIntyre and Faulk¹⁰ had isolated a glycoprotein from amnion and credited it to be responsible for suppressing any "foreign body" type reaction by acting on lymphocytes and preventing lymphoblastogenesis.

Amniotic membrane when used as an allograft in peritoneal cavity¹¹ or buried under skin⁴ has shown long term survival with no evidence of any immune reaction. Like wise, when used as xenograft from human to animals¹² or cattle to humans¹³ no significant antigenicity is revealed. Robson, Samburg and Krizek¹⁴ preferred to use the unseparated amniochorionic membrane with the chorion placed against the wound in deep second degree skin injuries. Although vascularization and vigorous "rejection" develops from the mesenchymal side of the chorion, both the procedures seem to be less intense when the epithelial cells are placed on the wound.

Walker¹⁵ has seen very little difference after 24 hours in the appearance of the membrane or of the wound whether amnion or chorion is placed next to it.

The difference in immunology of amnion and chorion is related to the presence of fragments of maternal decidua on the chorion.¹⁶ Animal experimentation suggests that the chorion alone can be responsible for the reaction. If the chorion is separated from the amnion and the mesenchymal side is applied to the host, a clear immune reaction may be observed.¹⁴

Functions of Amniotic Membrane

When used as a biological dressing, amniotic membrane has been credited with the following functions:

Vapour Barrier

By acting as an effective vapor barrier it prevents the evaporation of fluids from burn wound, thereby reducing insensible loss and in turn the overall fluid requirement of the body. Furthermore, by preventing evaporation from the wound surface, the temperature regulation mechanism is not over-strained and the caloric requirement to maintain body temperature is also correspondingly reduced.¹³ Vapor barrier property of amniotic membrane has been attributed to a firm bond between it and the wound, composed mostly of fibrin and elastin.¹⁷

When compared to homograft skin and porcine graft skin, amnion causes a very minor reduction of evaporation.¹⁸ Salisbury, Carnes and Enterline¹⁹ showed that in both full thickness and partial thickness wounds, allograft was as effective as sheet porcine skin and five times as effective as meshed porcine skin or amnion.

Bacteriostatic Function

This function is said to be due to the presence of antibodies, possibly allantoin, a bactericidal product of purine metabolism and lysozyme, a bacteriolytic protein.¹⁷

Adherence of amnion to the burn wound by eliminating its exposed status may itself lower bacterial count in the wound.^{12, 14, 20} The close adherence of the membrane to the wound is said to be via a fibrin and elastin.¹³

Furthermore, the amniotic membrane has a high thrombin activity which allows a very rapid and efficient attachment to living dermis or granulating tissue.²¹ This close adherence allows restoration of lymphatic integrity, protects circulating phagocytes from exposure and allows removal of surface debris and bacteria.²² It has been shown that amniotic membrane takes onto a granulating wound.⁴ This initial neo-vascularization is held responsible for effective decrease in bacterial counts.²³

Reduction of Pain

This is a frequently observed and well recognized quality of amniotic membrane when used as a skin substitute.²⁴⁻²⁶ It appears to follow diminished inflammation and possibly better state of hydration of wound bed.²⁷ Another explanation is that soft mucoid lining of amniotic membrane protects the exposed nerve endings from external irritant, the most important of which is the surrounding air.²⁸

Enhanced Wound Healing

Many workers have noted enhanced healing of wounds with the application of amniotic membrane.²⁹⁻³² Robson, Krizek, Koss and Samburg³³ in 1973 have subjectively noted rapidity of ingrowth of epithelium from the borders of the wound in full thickness defects and rate of re-epithelization of partial thickness burns appear to be increased by the use of amniotic membrane.^{26, 34-37}

Mechanism of healing by Amniotic Membrane

The most striking effect noted by Faulk et al³⁵ using amniotic membrane on chronic leg ulcer was the development of new vessels which they thought was due to some angiogenic factors acting on capillary endothelium.

Burgos³⁸ confirmed the presence of angiogenic and mitogenic factors in amniotic membrane and held them responsible for producing healing in the wounds. Freeze dried (lyophilized) amniotic membrane was used by Unger³⁹ on split skin graft donor sites who found it to be equivalent to an ordinary dressing and fail to notice any enhanced rate of healing.

The difference in potential of stimulating neo-vascularization and re-epithelization of fresh and lyophilized membrane is still controversial.⁴⁰ Klen and Skalska⁴¹ compared freeze dried amniotic membrane with freeze dried dermo-epidermal graft and concluded that chorion-amnion grafts were as effective as dermo-epidermal grafts.

Preparation of Amniotic Membrane

For clinical use the membrane can be prepared in the following forms: Fresh membrane as already described is obtained from the placenta at the time of delivery, either vaginal or caesarian section. Robson and Krizek¹² rinsed the membrane in a 0.025% solution of sodium hypochlorite and stored at 4°C in sterile solution containing penicillin. They showed that membranes remained sterile upto 6 weeks.

Dinno and associates²⁸ performed cultures to study sterilization of amniotic membranes. Preservation with 1:40 dilution of sodium hypochlorite revealed no positive cultures until 30 days.

Similar results were obtained with aqueous penicillin 50,000 units and streptomycin 1gm. in 400 ml. of normal saline. When kanamycin sulfate, 1.0 gm. in 400 ml. normal saline was used no positive cultures were found even at the end of 30 days.

Dried Membrane

Rao and Chandreshkaram¹³ after cleaning and rinsing the membranes spread them on a plastic sheet and allowed to dry in the open air. He found it to be equally effective when compared with the fresh.

Frozen Membrane

Amniotic membrane is frozen by passing through liquid nitrogen at -196°F. Cooling preserves the membrane for an indefinite time, produces bacteriologically pure and immunologically almost inert material.^{42,43}

Freeze Dried - Irradiated (Lyophilized)

In this process, membrane, after obtained from placenta is freeze dried at -60°C under vacuum (atmospheric pressure 102) for 48 hours. It is then irradiated with 2.5 mega rads (25 K Gray) in a batch type cobalt-60 irradiator.^{44,45} By the method of freeze drying there is sublimation of liquid moisture of membrane to gaseous state without having undergone the intermediate solid stage. This method helps the membrane to maintain its original size and shape with minimum cell rupture.⁴⁶ The freeze dried membrane can be readied for use by soaking it in normal saline for 1 minute.

Stabilized Amniotic Membrane

The idea of glutaraldehyde fixation of amniotic membrane was popularized which has led to the development of stabilized amniotic membrane (SAM).⁴⁷ Glutaraldehyde treatment required neither the antibiotics nor the use of special storage techniques and renders the amnion sterile as well as non-immunogenic. Successful use of glutaraldehyde treated amnion (SAM) is employed as a microvascular interpositional graft in experimental animals.⁴⁸

Storage of Amniotic Membrane

Dino³⁶ suggested the idea of human amnion bank which was seconded by Rao and Chandrasekhram¹³ also adding bovine amnions. Burgos and Faulk⁴⁰ describe a method to keep the amniotic membrane in culture for 2-3 weeks 50-90% viability.

Side of the Membrane towards Wound

There is great controversy as to which side of the membrane, amniotic or chorionic, should be applied next to the wound. Trelford⁴⁹ transplanted "trophoblastic tissue" (i.e. chorion) in sheep as an autograft and has shown that an immunological response occurs suggesting maternal decidua fragments may inadvertently be accompanying the chorion. He recommended that the mesenchymal side should be placed towards the wound for better survival. He had demonstrated that capillary and cellular invasion does not occur in the absence of chorion. Robson²³ considered "vascular invasion" to be the sole criteria for a dressing to be labeled as physiological and recommends that chorionic surface must be applied to deep or second degree burns to have any benefit. If amnion and chorion are separated and the amnion's mesenchymal side is applied to the host tissue, then vascularization and rejection phenomenon are not seen.⁵⁰ Walker⁵¹ saw very little difference after 24-48 hours in the appearance of the membrane or of the burn wound whether amnion or chorion was applied.

Method of Use

Before the membrane is applied, the wound should be prepared as for any dressing or for skin grafting. Surgical scrub with antiseptic and minimal debridement are followed by moist compression until oozing has stopped and the wound surface is reasonably dry. This procedure is preferably done in a clean sterile dressing room, observing all aseptic measures. No local or general anesthesia is required.

Membrane is applied with rough (chorionic) surface next to the wound. Care is taken to ensure no trapping of air bubbles between membrane and wound by gentle pressing. Membrane is followed by a layer of anti-bacterial gauze (e.g. Soframycin tulle), some moist gauze, dry gauze, cotton and bandage. Dressing should be changed along with the membrane at least every 48 hours and preferably after every 24 hours. Dressing should be continued for 7-10 days or until wound appears clinically clean. Split skin grafting should be done after 7-10 days or when wound contains less than 10⁵ organisms/gram of tissue.

Freeze dried irradiated membrane is also used as described above, but before application it is soaked in sterile saline for 1-2 minutes.

Use of Amniotic Membrane in other Surgical Disciplines

Apart from established use of amniotic membrane in acute superficial burn wounds and acute second-degree injuries, in

particular facial burns, some of the other indications are as follows:

Following facial dermabrasion⁵², vaginal reconstruction^{53,54}, replacing nasal mucosa⁵⁵, bladder wall reconstruction⁵⁶, Stevens-Johnson syndrome⁵⁷, non-healing leg ulcers^{37,58}, reconstruction of the floor of the mouth following total glossectomy⁵⁹, micro-vascular interpositional grafts⁴⁸, conjunctival² and corneal defects⁶⁰, graft donor sites²⁰ and radiation burns.⁶¹

Advantages of Amniotic Membrane as a Biological Membrane

It is readily available at no cost if fresh²⁷, sterilization, storage and application are simple, prevents fluids, protein and energy loss⁶², combats infection⁶³, promotes healing⁶⁴ and relieves pain.²⁶

Disadvantages

It is highly fragile and becomes firmly adherent to the wound. Attempts to remove it, even after soaking the area, can cause considerable bleeding and pain to the patient.

Amniotic Membrane in a Developing Country

Considering the properties of amniotic membrane and the easy availability, low cost of procurement and cheap storage makes it appear to be a useful dressing material for burn wounds and other non healing skin lesions in developing countries.^{26,61,65-67}

References

1. Walker AB. Use of amniotic membrane for burn wound coverage in Wise DL (ed.) Burn wound coverings. Boca Raton, CRC Press, 1984, p. 57.
2. DeRoth A. Plastic repair of conjunctival defects with fetal membranes. Arch Ophthalmol 1940;23 522.
3. Troensegaard-Hansen E. Amniotic grafts in chorionic skin ulceration. Lancet 1950;1: 859.
4. Douglas B, Conway H, Stark RB, et al. The fate of homologous and heterologous chorionic transplants as observed by the transparent tissue chamber technique in the mouse. Plas Reconstr Surg 1954;13:125.
5. Sterling JA. Use of amniotic membranes to cover surface defects due to flame burns. Am J Surg 1956;91:940.
6. Troensegaard-Hansen E. Amnion implantation in peripheral vascular disease. Br Med J 1956;4:262.
7. Pigeon J. Treatment of second degree burns with amniotic membranes. Can Med Assoc J 1960;83 844.
8. Dino BR, Eufemio GG, DeVilla MS, et al. The use of fetal membrane homografts in the local management of burns. J Philippine Med Assoc 1965;41: 890.
9. Schwarts AL, Forster CS and Liggins GC. Human amnion metabolism. Am J Obstet Gynecol 1977;127: 470.
10. McIntyre JA and Faulk WP. Antigens of human trophoblasts: effect of heterologous and anti-trophoblast sera on lymphocyte response in utero. J Exp Med 1979;149: 824.
11. Trelford JD, Hanson FW, Anderson DG, et al. Implanted amniotic membranes as an autograft and as an allograft. J Med 1975;6:169.
12. Robson MC and Krizek TJ. The effect of human amniotic membranes on the bacterial population of infected rat burns. Ann Surg 1973;177:144.
13. Rao TV and Chandrasekhram V. Use of dry human and bovine amnion as a biological dressing. Arch Surg 1981;116: 891.

14. Robson MC, Samburg JL, Krizek TJ. Quantitative comparison of biological dressings. *J Surg Res* 1973;14: 431.
15. Walker AB. Use of amniotic membranes for burn wound coverage in Wise DL (ed.) *Burn Wound coverings*, CRC Press Boca Raton, 1984, p. 58.
16. Kelton PL: Principles of skin grafts in Kelton PL (ed.) in *Selected Readings in Plastic Surgery 4/2*: Dallas, Texas: 1986 pp. 10-11.
17. Walker AB, Cooney DR and Allen JE. Use of fresh amnion as a burn dressing. *J Pediatr Surg* 1977;12:391.
18. Lamke LO. The influence of different skin grafts on the evaporative water loss from burns. *Scand Plast Reconstr Surg* 1997;5:82.
19. Salisbury RE, Carnes RW, Enterline D: Biological dressings and evaporative water loss from burn wounds. *Ann Plast Surg* 1980;5/4:270.
20. Quinby WC, Hoover HC, Schefflan M, et al. Clinical trials of amniotic membranes in burn wound care. *Plast Reconstr Surg* 1982;70:711.
21. Walker AB. Use of amniotic membranes for wound coverage in Wise DL (ed.) *Burn wound coverings*, Boca Raton: CRC Press, 1984, p. 60.
22. Burleson R, Eiseman B. Mechanisms of antibacterial effect of biological dressings. *Ann Surg* 1973;117:181.
23. Robson MC. Invited editorial comment. In: Rao TV and Chandrasekharam V. Use of human and bovine amnion as a biological dressing. *Arch Surg* 1981;116: 891.
24. Sharma SC, Bagree MM, Bhat et al. Amniotic membrane is an effective burn dressing material. *Jpn J Surg* 1985;15/2:140.
25. Landi G, Fortuna A, Mengozzi E, et al. Medication burns with amniotic membrane. *G Ital Dermatol/Minerva Dermatol* 1977;112:343.
26. Kasi N, Durrani KM, Siddiqui MA. Human amniotic membrane as a versatile biological dressing. A preliminary report. *J Pak Med Assoc* 1987;37:290.
27. Rao TV, Chandrasekharam V. Human amnion as a dressing material in burns. *Indian J Surg* 1981;43:561.
28. Dino BR, Eufemio GG, DeVilla MS. Human amnion: the establishment of an amnion bank and its practical application in surgery. *J Philippine Med Assoc* 1966;42:230.
29. Trelford JD, Anderson DG, Hanson FW, et al. Wound healing and the amniotic membrane. *J Med* 1975;6:383.
30. Eldad A, Stark M, Anais D, et al. Amniotic membrane as a biological dressing. *S Afr Med J* 1977;51:272.
31. Check WA: Encouraging news on temporary coverings for wounds, reporting on exhibit by Walker AB at American College of Surgeons, Atlanta, 1979. *JAMA* 1980;244:2493.
32. Androulakis I, Petrochilos I, Kalfarentzos P, et al. The effect of amniotic membrane on the microcirculation of burns (preliminary report) *Excerpta Medica I.C.S.*, 1977;409 973.
33. Robson MC, Krizek TJ, Koss N, et al. Amniotic membranes as a temporary wound dressing. *Surg Gyne Obstet* 1973;136:904.
34. Colocho G, Graham WP, Greene AE, et al. Human amniotic membranes as a physiological wound dressing. *Arch Surg* 1974;109: 370.
35. Faulk WP, Matthews RN, Stevens PJ, et al. Human amnion as an adjunct in wound healing. *Lancet* 1980;1:1156.
36. Matthews RN, Bennett JP, Faulk WP. Wound healing using amniotic membranes. *Br J Plast Surg* 1981;34:76.

37. Bennett JP, Matthews RN, Faulk WP. Treatment of chronic ulceration of the legs with human amnion. *Lancet* 1980;1:1153.
38. Burgos H. Angiogenic and growth factors human amnio-chorion and placenta. *Europ J Clin Invest* 1983;13:289.
39. Unger MG, Roberts M. Lyophilized amniotic membrane on graft donor site. *Br J Plast Surg* 1976;29:99.
40. Burgos H & Faulk WP. The maintenance of human amniotic membrane in culture. *Br J Obstet Gynecol* 1981;88:294.
41. Klen R, Shalska H. A comparison of dermo epidermal and chorion amniotic grafts in the treatment of burns. *Acta Chir plast* 1976;18:225.
42. Vitale R, Iaia A, Sferrazza G, et al. A biological dressing for burn wound. *Ita Riv Ital Chir Plast* 1981;13 127.
43. Lorusso P, Geraci V, Masellis M. The treatment of superficial burns with biological and synthetic material. Frozen amnion and Biobrane. *Ann Burn Med Club* 1989;2:79.
44. Siddiqui MA. Freeze-dried, radiated sterilized human amniotic membrane as a biological dressing for burns and chronic ulcers. Liaquat Medical College, Jamshoro 1981 (unpublished report).
45. Notea E, Hirshowitz B, Karev A, et al. Lyophilized amnion in burns and skin loss. *Harefuah* 1975;16: 265.
46. Nazri MM. Freeze-drying: the latest in food technology. *Friday Dawn Magazine*, Karachi, February 1990, 16: II.
47. Thomson PD, Parks DH. Amnion as a wound dressing in Wise DL (ed.) *Burn wound coverings*, Boca Raton: CRC Press, 1984, pp. 48-49.
48. Gray KJ, Shenaq SM, Engelmann UH, et al. Use of amnion for microvascular interpositional grafts. *Plast Reconstr Surg* 1986;79:778.
49. Trelford JD, Anderson DG, Hanson FW, et al. Consideration of the amnion as an autograft and as an allograft in sheep. A preliminary report. *J Med* 1972;3:231.
50. Trelford JD, Hanson FW, Anderson DG, et al. Amnion autografts, permanent structure. *J Med* 1975;6:243.
51. Walker AB. Use of amniotic membranes for burn wound coverage in Wise DL (ed.) *Burn wound coverings*, Boca Raton: CRC Press, 1984, p. 62.
52. Kucan JO, Robson MC, Parsons RW. Amniotic membrane as dressing following facial dermabrasion. *Ann Plast Surg* 1982;8:523.
53. Tancer ML, Katz M, Veridiano NP. Vaginal epithelization with human amnion. *Obstet Gynecol* 1979;54:345.
54. Tozum R. Homotransplantation of amniotic membrane for the treatment of congenital absence of vagina. *Gynecol Obstet* 1977;14:553.
55. Zohar Y, Talmi YP, Frankelstein Y, et al. Use of human amniotic membrane in oto-laryngologic practice. *Laryngoscope* 1987;97:978.
56. Fishman IJ, Flores FN, Scott FB, et al. Use of fresh placental membranes for bladder reconstruction. *J Urology* 1987;138: 1291.
57. Prasad JK, Feller I, Thomson PD. Use of amnion for the treatment of Stevens-Johnson Syndrome. *J Trauma* 1986;26: 945.
58. Ward DJ, Bennett JP. The long term result of the use of human amnion in the treatment of leg ulcers. *Br J Plast Surg* 1984;37:191.

59. Bapat CV, Kothary PM. Preliminary report on acceleration of wound healing by amnion membrane graft. *Indian J Med Res* 1974;62:1342.
60. Ucakhan OO, Koklu G, Firat E. Non-reserved human amniotic membrane transplantation in acute and chronic chemical injuries. *Cornea* 2002;21:169-72.
61. Walker AB. Use of amniotic membranes for burn wound coverage in Wise DL (ed.) *Burn wound coverings*, Boca Raton: CRC Press,, 1984, p. 69.
62. Chuntrasakul C. Clinical experience with the use of amniotic membranes as a temporary dressing in the treatment of burns and other surgical open wounds. *Med Assoc Thailand J* 1977;60:66.
63. Robson MC, Krizek TJ. Clinical experiences with amniotic membranes as a temporary biological dressing. *Conn Med* 1974;38:449.
64. Bose B. Burn wound dressing with human amniotic membrane. *Ann Royal Coll Surg Engl* 1979;61:444.
65. Piserchia NE, Akenzua GI. Amniotic membrane dressing for burns in children: a cheap method of treatment for developing countries. *Trop Geogr Med* 1981;33: 235.
66. Ramakrishnan KM, Rao DK. Human amniotic membrane as a temporary dressing in complicated burns in a developing country. *J Burn Care Rehabil* 1984;4:202.
67. Ramakrishnan KM, Jayaraman V. Management of partial thickness burn wounds by amniotic membrane: a cost effective treatment in developing countries. *Burns* 1997; 23:33-6.