

Assessment of risk factor profile in young patients undergoing elective coronary artery bypass grafting surgery in Armed Forces Institute of Cardiology/National Institute of Heart Disease, a tertiary care cardiac facility

Mishal Iqbal¹, Mehak Iqbal², Muhammad Afsheen Iqbal³, Farrah Pervaiz⁴, Hamza Jamil⁵, Syed Hashim Ali Inam⁶

Abstract

Objective: To identify modifiable and non-modifiable risk factors among young patients undergoing elective coronary artery bypass graft surgery.

Methods: The descriptive cross-sectional study was conducted at the Department of Adult Cardiac Surgery of the Armed Forces Institute of Cardiology and National Institute of Heart Disease, Rawalpindi, Pakistan, from January 1, 2015, to April 24, 2018, and comprised patients who underwent elective coronary artery bypass grafting. Data was collected using a predesigned questionnaire, and the risk factors studied were hypertension, diabetes, smoking, serum cholesterol, family history and body mass index. Data was analysed using SPSS 23.

Results: Of the 1270 patients, 838(66.1%) were males and 432(33.9%) were females. The overall mean age was 39.29±8.64 years (range: 18-49 years). Of the total, 319(25.2%) patients were hypertensive; 170(13.3%) were diabetics; 303(23.9%) had history of smoking; 566(44.6%) were obese.

Conclusion: There was a high prevalence of obesity, hypertension and smoking in the patients undergoing coronary artery bypass graft surgery at age <50 years.

Keywords: Coronary artery disease, Risk factors, Young patients. (JPMA 71: 98; 2021)

DOI: <https://doi.org/10.47391/JPMA.712>

Introduction

The incidence of coronary artery disease (CAD) at age <45 years is characterised as young CAD.¹ However, numerous studies have deliberated the age limit varying from 35 years to 55 years in the spectrum of young CAD.²⁻⁴ This field of cardiology has acquired significance recently due to higher prevalence in this age group over the last few decades, with fluctuating risk factor profiles and alteration in prognosis as well as longevity after an acute coronary episode.

Prevalence of conventional risk factors like hypertension (HTN), diabetes mellitus (DM), smoking, dyslipidaemia and obesity accounts for about 85% to 90% of premature CAD patients. Often young CAD patients have multiple concomitant risk factors contributing to the disease.^{5,6} The most common risk factor associated with young CAD seems to be smoking. The prevalence of smoking in individuals aged <45 years, with CAD, was documented as 60-90% compared to 24-56% in individuals aged >45 years.⁷ Smoking in the presence of additional risk factors, like DM, HTN and obesity, predisposes a young individual

to increased risk of future acute coronary events.^{7,8}

The current study was planned to identify modifiable and non-modifiable risk factors among young patients undergoing elective coronary artery bypass graft (CABG) surgery.

Patients and Methods

The descriptive cross-sectional study was conducted at the Armed Forces Institute of Cardiology and the National Institute of Heart Diseases (AFIC/NIHD) from January 1, 2015, to April 24, 2018, and comprised patients aged 18-49 years who underwent elective CABG. After approval from the institutional ethical review board, the sample size was calculated with 29% prevalence noted in statistics.⁹ To increase the validity of results the sample size was increased over four-fold. Patients who had undergone emergency CABG were excluded. A questionnaire was designed to collect data from the patients that included history and biochemical markers.

HTN, DM, smoking, serum cholesterol, family history of cardiovascular disease, and body mass index (BMI) were studied as the risk factors in patients. HTN was defined as systolic blood pressure (SBP) >140mmHg and diastolic blood pressure (DBP) 90 mmHg, as per the World Health Organisation (WHO) criteria¹⁰ and was further categorised into two groups controlled with medication and uncontrolled HTN. DM was defined as random blood

¹Department of Medicine, Pak Emirates Military Hospital, Rawalpindi, Pakistan; ^{2,5,6}Department of Medicine, Army Medical College, Rawalpindi, Pakistan; ³Department of Cardiac Surgery, Armed Forces Institute of Cardiology, National Institute of Heart Diseases, Rawalpindi, Pakistan; ⁴Department of Research and Development, Armed Forces Institute of Cardiology, National Institute of Heart Diseases, Rawalpindi, Pakistan.

Correspondence: Misha Iqbal e-mail: mi8@hotmail.co.uk

glucose (RBG) >200mg/dl or fasting blood glucose (FBG) >126mg/dl.¹¹ Diabetics were divided into three sub-groups: those on oral therapy, on insulin, and patients whose diabetes was controlled by diet. On the basis of smoking history, the subjects were classified into three categories; current smokers, past smokers who had given up smoking >8 weeks ago, and non-smokers who had never smoked. Serum cholesterol level was divided into three ranges; normal, >200 and level >300. Patients were asked about their family history regarding CAD. Body mass index (BMI) was calculated by dividing the patients' weight (in kg) by the patients' height squared (in m²). BMI values were categorized into normal (18.0-22.9 kg/m²), overweight (23-24.9 kg/m²) and obese (>25 kg/m²).¹²

Data was analysed using SPSS 23. Continuous variables were expressed as mean±standard deviation (SD) and qualitative variables were expressed as frequencies and percentages.

Results

Of the 1270 patients, 838(66.1%) were males and 432(33.9%) were females. The overall mean age was 39.29±8.64 years. Of the total, 319(25.2%) patients were hypertensive, 170(13.3%) were diabetics and 303(23.9%) had history of smoking. The cholesterol level was recorded

for 129 patients and the mean level was 160±45.9. Of these patients, 21(16.2%) had Cholesterol level >200mg/dl. Family history of cardiovascular disease was known in 221(17.4%) patients and 6(0.5%) had no knowledge about their family history. The mean weight was 67.53±15.24kg, the mean height was 165.21±11.04cm, and the mean BMI was 24.66±5.36 kg/m². Overall, 566 (44.6%) patients were obese (Table).

Discussion

The current study looked into six risk factors for ischaemic heart disease (IHD) in patients aged 49 years and younger. IHD was more prevalent among the males. The most common factor present in the study population was a raised BMI, followed by HTN, smoking, family history and DM. DM was the least common risk factor.

Conventionally, coronary heart disease is considered a disease of the male gender, but it is becoming common in women and a higher prevalence of HTN, DM and obesity in older women has led to an increase in the number of women suffering from coronary heart disease.¹³ Smoking and lipid abnormalities in women also play a major role.¹³ IHD is the most common cause of death among women in western countries.¹⁴ Another study showed that heart disease develops 7-10 years later in women than in men.¹⁵

In the current study, 319 (25.2%) patients suffered from HTN. A study in India stated 70.9% hypertensives.¹⁶ A study in Iran on CAD patients aged <45 years showed 37% HTN.¹⁷ In a Karachi-based CAD analysis, only 19% patients were hypertensive¹⁸ which was lower than our findings.

The current study had 170 (13.3%) diabetics. A study showed DM incidence of 10% in CAD patients aged <40 years.¹⁹ A study in India reported a higher DM prevalence of 47.5%, but the mean age of patients in that study was 59.73 years.¹⁶ In Iran, it was 27%,¹⁷ and in Karachi it was 15.4%.¹⁸ In the current study, 303 (23.9%) patients were smokers compared to 58.7% reported earlier.² Foroughi M et al. documented 54% prevalence among Iranians.¹⁷ The prevalence of smoking in young CAD patients was 92% in another study and smoking was more common among younger patients.²⁰ In a study done on risk factors for premature CAD, the prevalence of smoking was higher in patients aged <40(58.7%) years compared to those aged >90 years (43%).²¹

As for hyperlipidaemia, it was 23(17.7%) in the current study compared to 39% in Iranian patients.¹⁷ A study in Seattle reported that 31% patients <90 years had hyperlipidaemia, and the lipid abnormalities were more common among males <40 and females <50 years.²²

In the current study, 221(17.4%) patients had a family

Table: Demographic characteristics (n= 1270).

S. No	Characteristic	n (%)
1	Hypertension	319 (25.2)
	Controlled (medication)	312 (24.6)
	Uncontrolled	7 (0.6)
2	Diabetes	170 (13.3)
	Oral Therapy	130 (10.2)
	Insulin	30 (2.4)
	Dietary Control	10 (0.8)
3	Smoking	303 (23.9)
	Current smokers	111 (8.7)
	Past smokers	192 (15.1)
4	Cholesterol	
	Normal	106 (82.3)
	Above normal	23 (17.7)
	>200	21 (16.27)
	>300	2 (1.55)
5	Family History of Coronary Artery Disease	
	Known	221 (17.4)
	Unknown	1049 (82.6)
6	Weight	67.53±15.24
7	Height	165.21±11.04
8	Body Mass Index (BMI)	
	Underweight	117 (9.2)
	Normal	362 (28.5)
	Overweight	225 (17.7)
	Obese	566 (44.6)

history of cardiovascular disease, whereas it was 38% in Iran¹⁷ and 39% in a study done in London. Positive family history has been reported to be more common among young men.^{17,23} Christus T et al. documented a positive family history of CAD among younger patients (39%) compared to older patients (11%).⁴

Mean BMI in the study was $24.9912 \pm 5.39 \text{ kg/m}^2$ whereas it was $28 \pm 4.7 \text{ kg/m}^2$ in Iran.¹⁷ Raised BMI was the most prevalent 791 (62.3%) risk factor in our study in which 225 (17.7%) patients were overweight and 566 (44.6%) were obese. In a study in Karachi 24% subjects were obese.¹⁸

The current study has limitations. It was done at a single centre and the cholesterol levels were recorded for only a portion of the sample. Also, the study population was limited to elective patients only, and majority belonged to a poor socioeconomic background and were mainly uneducated. Thus, they were unsure of their family history.

To calculate the true incidence and prevalence of these risk factors, future studies are required to evaluate the epidemiological transition toward a rising trend of CAD in young population and to guide evidence-based practices in prevention and screening of high-risk individuals.

Conclusion

There was a high prevalence of obesity as well as non-modifiable risk factors, like hypertension, in patients aged <50 years undergoing CABG surgery.

Disclaimer: None.

Conflict of interest: None.

Source of Funding: None.

References

- Egred M, Viswanathan G, Davis G. Myocardial infarction in young adults. *Postgrad Med J*. 2005;81:741-5.
- Konishi H, Miyauchi K, Kasai T, Tsuboi S, Ogita M, Naito R, et al. Long-term prognosis and clinical characteristics of young adults (≤ 40 years old) who underwent percutaneous coronary intervention. *J. Cardiol*. 2014;64:171-4.
- Gupta R, Misra A, Vikram NK, Kondal D, Gupta SS, Agrawal A, et al. Younger age of escalation of cardiovascular risk factors in Asian Indian subjects. *BMC Cardiovasc. Disord*. 2009;9:28.
- Christus T, Shukkur A, Rashdan I, Koshy T, Alanbaei M, Zubaid M, et al. Coronary artery disease in patients aged 35 or less—a different beast? *Heart views*. 2011;12:7.
- Khot UN, Khot MB, Bajzer CT, Sapp SK, Ohman EM, Brener SJ, et al. Prevalence of conventional risk factors in patients with coronary heart disease. *JAMA*. 2003;290:898-904.
- Celik T, Ilyisoy A. Premature coronary artery disease in young patients: an uncommon but growing entity. *Int. J. Cardiol*. 2010;144:131-2.
- Genest JJ, McNamara JR, Salem DN, Schaefer EJ. Prevalence of risk factors in men with premature coronary artery disease. *Am J Cardiol*. 1991;67:1185-9.
- Pineda J, Marín F, Marco P, Roldán V, Valencia J, Ruiz-Nodar JM, et al. Premature coronary artery disease in young (age < 45) subjects: interactions of lipid profile, thrombophilic and haemostatic markers. *Int. J. Cardiol*. 2009;136:222-5.
- Health statistics of Pakistan. WHO NCD Country Profile. [Online] [Cited 2018 May 10]. Available from: URL:https://www.who.int/nmh/countries/pak_en.pdf
- Organization WH. Hypertension control: report of a WHO Expert Committee. 1996.
- Kerner W, Brückel J. Definition, classification and diagnosis of diabetes mellitus. *Exp Clin Endocrinol Diabetes*. 2014;122:384-6.
- WHO. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies [Internet] Available from: https://www.who.int/nutrition/publications/bmi_asia_strategies.pdf [Cited 05 December 2020]
- Tan YY, Gast G-CM, van der Schouw YT. Gender differences in risk factors for coronary heart disease. *Maturitas*. 2010;65:149-60.
- Abbey SE, Stewart DE. Gender and psychosomatic aspects of ischemic heart disease. *J Psychosom Res*. 2000;48:417-23.
- Maas A, Appelman Y. Gender differences in coronary heart disease. *NETH HEART J.* 2010;18:598-603.
- Kasliwal R, Kulshreshtha A, Agrawal S, Bansal M, Trehan N. Prevalence of cardiovascular risk factors in Indian patients undergoing coronary artery bypass surgery. *JAPI*. 2006;54:371-5.
- Foroughi M, Ahranjani S, Ebrahimian M, Saieedi M, Safi M, Abtahian Z. Coronary artery disease in Iranian young adults, similarities and differences. *Open J Epidemiol*. 2014;4:19-24.
- Iqbal S, Dodani S, Qureshi R. Risk factors and behaviours for coronary artery disease (CAD) among ambulatory Pakistanis. *J Pak Med Assoc*. 2004;54:261-266
- Cole JH, Miller JI, Sperling LS, Weintraub WS. Long-term follow-up of coronary artery disease presenting in young adults. *J Am Coll Cardiol*. 2003;41:521-8.
- Zimmerman FH, Cameron A, Fisher LD, Grace N. Myocardial infarction in young adults: angiographic characterization, risk factors and prognosis (Coronary Artery Surgery Study Registry). *J Am Coll Cardiol*. 1995;26:654-61.
- Mukherjee D, Hsu A, Moliterno DJ, Lincoff AM, Goormastic M, Topol EJ. Risk factors for premature coronary artery disease and determinants of adverse outcomes after revascularization in patients ≤ 40 years old. *Am J Cardiol*. 2003;92:1465-7.
- Goldstein JL, Hazzard WR, Schrott HG, Bierman EL, Motulsky AG. Hyperlipidemia in coronary heart disease I. Lipid levels in 500 survivors of myocardial infarction. *J Clin Invest*. 1973;52:1533-43.
- Chen L, Chester M, Kaski JC. Clinical factors and angiographic features associated with premature coronary artery disease. *Chest*. 1995;108:364-9.