

Improvement in knowledge of healthcare professionals attending Neonatal Life Support (NeoLiS) training workshop

Zeeshan Munir¹, Sikandar Hayat², Bushra Fatima³, Muhammad Anwar⁴, Mazhar Qadir⁵, Irfan Waheed⁶

Abstract

Objective: To assess the improvement in knowledge of healthcare professionals after attending Neonatal Life Support training workshop.

Methods: The interventional study with secondary data was conducted in the Neonatology Department of the Children's Hospital and the Institute of Child Health, Lahore, Pakistan, using 2-year data of all the training workshops from 2017 to 2018. The participants were consultants, post-graduate residents, nurses, lady health visitors, lady health workers and rescue workers. Knowledge was tested through a true/false questionnaire regarding resuscitation skills both before and after the hands-on workshop. The score of each candidate was recorded on a results sheet. Baselines scores were compared with respective post-workshop test. Data was analysed using SPSS 20.

Results: Of the 52 workshops, 35(%) were conducted in 2017 and 17(%) in 2018, and of the 1350 participants, 1080(81%) were doctors, 257(18%) were nurses and 13(1%) were paramedics and rescue workers. Mean overall baseline score was 71.96 ± 8.38 compared to 86.60 ± 5.48 post-intervention ($p < 0.001$).

Conclusion: The Neonatal Life Support training workshop resulted in significant gain in knowledge about neonatal resuscitation by healthcare professionals.

Keywords: Neonatal resuscitation, Neonatal training workshops, Helping babies breathe, Neonatal life support.

(JPMA 71: 1996; 2021) DOI: <https://doi.org/10.47391/JPMA.04-526>

Introduction

About 5.4 million children aged <5 years die worldwide every year, and 45% of them are newborns. According to the World Health Organisation (WHO), about two-third of these newborns die in the first 24 hours. Pakistan ranks among the countries with highest neonatal mortality rate (NMR). In 2017, the NMR in Pakistan was estimated to be 42/1000 live-births.^{1,2}

The causes include healthcare budgetary constraints, low rates of exclusive breastfeeding, poor female literacy rates, lack of antenatal visits, low institutional delivery rates and poor access to skilled birth attendants (SBAs).^{2,3}

Birth-related events, like asphyxia neonatorum, represent one of the three most common causes of global neonatal mortality which is estimated to cause 0.7 to 1 million deaths per year globally, and most of these deaths occur in the developing countries. In Pakistan, 298,000 neonates die annually, and asphyxia neonatorum accounts for 23% of these deaths.^{4,5}

The initiation of breathing is a critical step in the physiological transition of a baby from intra-uterine to

extra-uterine life. Approximately 85% of babies born at term will initiate spontaneous breathing within 10-30 seconds of birth, an additional 10% will respond to drying and stimulation, 3% will initiate respirations after positive-pressure ventilation (PPV), 2% will be intubated to support respiratory function, and 0.1% will require chest compressions (CCs) and/or drugs to achieve this transition.^{6,7} Effective resuscitation during the Golden Minute, which is the first minute of life post-delivery, can improve mortality and morbidity from this preventable cause. Therefore, training programmes on neonatal resuscitation for all health workers involved in the management of the newborn at birth are vital to improve neonatal survival. In developed countries, different training programmes are conducted, like helping babies breathe (HBB), neonatal resuscitation programme (NRP) and newborn life support (NLS). All these training programmes are similar, but with minor differences and are based on the guidelines of the International Liaison Committee on Resuscitation (ILCOR). Such training programmes are regularly conducted and monitored by resuscitation councils, and studies have shown improvement in knowledge and skills of healthcare workers after attending these workshops.⁸⁻¹²

In Pakistan, a training programme called the Neonatal Life Support (NeoLiS) was developed in 2008 which was also based on ILCOR guidelines and endorsed by the Pakistan

^{1-3,5,6}Department of Neonatology, The Children's Hospital and The Institute of Child Health, Lahore, Pakistan; ⁴Department of Neonatology, Victoria Hospital, Bahawalpur, Pakistan.

Correspondence: Zeeshan Munir. e-mail: drzeeshanmunir@gmail.com

Annexure: NEONATAL LIFE SUPPORT WORKSHOP

Department of Neonatology,
The Children's Hospital & the Institute of Child Health, Ferozepur Road, Lahore
KNOWLEDGE CHECK
Time allowed: 15 minutes

Name: _____ Marks obtained: _____

Please read the statements and encircle the correct answer: - (T = True, F = False)



- | | | |
|--|---|---|
| 1. Every delivery should be attended by at least one skilled person. | T | F |
| 2. If chest is not rising with bag & mask ventilation, the seal between mask and face may be inadequate. | T | F |
| 3. The correct way to stimulate the baby is to slap the back. | T | F |
| 4. In the first minute after birth you should help the baby breathe. | T | F |
| 5. Umbilical cord of spontaneously crying baby should be clamped and cut immediately after birth. | T | F |
| 6. The correct depth of chest compression is 1/5th of antero-posterior diameter of the chest. | T | F |
| 7. A vigorously crying baby, delivered with meconium staining, needs immediate endotracheal intubation. | T | F |
| 8. Babies with suspected congenital diaphragmatic hernia should have continuous bag & mask ventilation. | T | F |
| 9. Overhead heat is more important than drying a newborn. | T | F |
| 10. During cardiopulmonary resuscitation, the lungs should be inflated at a rate of 30 breaths per minute. | T | F |
| 11. During resuscitation of a newborn, the ratio of breaths to chest compressions should be 1: 2. | T | F |
| 12. Hypothermia can make cardiopulmonary resuscitation difficult. | T | F |
| 13. Endotracheal intubation is essential for effective resuscitation. | T | F |
| 14. After a successful resuscitation, the notes shall not be written. | T | F |
| 15. The parents shall be informed whether the resuscitation is successful or not. | T | F |

Paediatric Association (PPA). The current study was planned to assess the improvement in knowledge of healthcare professionals after attending the NeOLiS training workshop.¹³

Subjects and Methods

The interventional study with secondary data was conducted in the Neonatology Department of the Children's Hospital and the Institute of Child Health, Lahore, Pakistan, using 2-year data of all the training workshops from 2017 to 2018. The workshop was developed keeping in view the facilities available for effective resuscitation at different levels of healthcare establishments in the country, and without compromising on ILCOR guidelines. The first part of technique related to the Golden Minute was influenced by HBB, but continued further on to the administration of drugs. Thus, it was simpler, evidence-based and avoided confusion in techniques in different resuscitation programmes developed by different countries. Moreover, it was easier for the facilitators to follow one algorithm where they taught only the initial part, up to airway management, to nurses and midwives working in level-I healthcare facilities, and continued further to the injection of drugs while training doctors and nurses in Level-III hospitals. The workshops were regularly arranged and monitored by the Neonatal Resuscitation Council of PPA and the Directorate of NeOLiS. Each workshop consisted of 25-30 participants who were doctors, both consultants and post-graduate residents,

nurses and paramedics, including lady health visitors (LHVs), lady health corks (LHWs) and rescue workers. All individuals who participated in the workshops were included, but those who did not completely fill the questionnaire were excluded. Knowledge was tested through a true/false questionnaire regarding resuscitation skills (Annexure) both before and after the hands-on workshop. Each question carried one mark. The score of each candidate was recorded on a results sheet. Baselines scores were compared with respective post-workshop test. Data was analysed using SPSS 20. $P < 0.05$ was considered significant.

Results

Of the 52 workshops, 35(%) were conducted in 2017 and 17(%) in 2018, and of the 1350 participants, 1080(81%) were doctors, 257(18%) were nurses and 13(1%) were paramedics and rescue workers. Mean overall baseline score was 71.96 ± 8.38 compared to 86.60 ± 5.48 post-intervention ($p < 0.001$) (Table 1).

The difference was significant for all the three categories of healthcare workers (Table 2).

Table-1: Results of all workshops.

	Mean±SD	p-value
Mean of all 52 Pre-workshop results	71.96±8.38	<0.01
Mean of all 52 Post-workshop results	86.60±5.48	

SD=Standard Deviation

Table-2: Results of all categories (n=52).

	Mean±SD	p-value
Mean of Pre-workshop results of doctors	73.41±8.48	<0.01
Mean of Post-workshop results of doctors	87.99±4.83	
Mean of Pre-workshop results of Nurses	66.84±10.49	<0.01
Mean of Post-workshop results of Nurses	82.74±6.26	
Mean of Pre-workshop results of Paramedics	52.26±10.14	<0.01
Mean of Post-workshop results of Paramedics	75.80±7.45	

SD=Standard Deviation

Discussion

Resuscitation at birth is the cornerstone to preventing birth asphyxia and associated morbidity, like cerebral palsy (CP). It is estimated that each CP patient consumes about £750,000 during his/her lifetime for treatment which

otherwise could be spent on national development projects. Preventing asphyxia, therefore, not only saves a newborn from lifelong disability, but also saves a family from perpetual emotional trauma and the nation from diverting huge amount of funds for treatment and rehabilitation of disabilities associated with CP. An effective neonatal resuscitation course is expected to improve the knowledge of all cadres of healthcare professionals.¹⁴

A study on HBB workshop assessed skills of physicians, nurses and other health workers, showing overall improvement from pre-workshop mean score of 8.7/10 to a post-workshop mean of 9.4/10.¹⁵ Another study assessed the educational effectiveness of HBB in healthcare professionals and found marked improvement in knowledge and skills after such courses.¹⁶ The results of these studies are consistent with current findings.

Healthcare professionals, like physicians, nurses and paramedics, have different sets of background knowledge regarding resuscitation and therefore, the outcome of these resuscitation workshops sometimes depend on their previous knowledge. A study showed that there was only modest gain in knowledge after a workshop as it had enrolled only undergraduate students who had no background knowledge of resuscitation.¹⁷ Similarly, a study on Paediatric residents also showed modest gain in knowledge after NRP.¹⁸ Another study on HBB included physicians and nurses, and reported marked improvement in knowledge post-workshop. However, the overall score percentage of nurses was lower than that of the doctors though post-training even those nurses who commonly did not perform these skills in real-life situations were also able to perform at a level similar to that of the physicians.¹⁹ The current study also showed significant improvement in knowledge of staff nurses and paramedics after attending NeoLiS workshop. However, the overall results of staff nurses and paramedics were lower compared to those of the doctors. This is likely due to the previous background core knowledge of doctors about resuscitation.

The knowledge and skills gained by healthcare professionals during neonatal resuscitation workshops are expected to decline over time and need to be updated on a regular basis. A study on NRP assessed knowledge using a questionnaire and concluded that the scores significantly improved after the course. However, scores at 6-month follow-up were significantly decreased from post-test, while remaining significantly higher compared to the pre-test performance.²⁰ Another study showed that skills declined more than knowledge over time.²¹ Likewise, in the current study, immediate assessment of knowledge after the workshop showed significant gains in results. However, further studies are required to assess the long-term impact

of these workshops on healthcare professionals.

Conclusion

The NeoLiS training workshops resulted in significant gain in knowledge about neonatal resuscitation by healthcare professionals when assessed immediately after the workshop.

Disclaimer: None.

Conflict of Interest: None.

Source of Funding: None.

References

1. WHO. Health-related Millennium Development Goals. World Health Statistics. [Online] 2018 [Cited 2021 Jan 15] Available from: URL: http://www.who.int/gho/publications/world_health_statistics/2018/en/
2. National institute of population studies (NIPS) [PAKISTAN] and ICF. Pakistan Demographic and Health survey 2017-18. Islamabad, Pakistan and Rockville, Maryland, USA: NIPS and ICF.
3. Khaliq F, Ahmed W. State of health sector of Pakistan. [Online] 2018 [Cited 2021 April 06] Available from: URL: <http://www.sbp.org.pk>
4. Umar LW, Ahmad H, Isah A, Idris HW, Hassan L, Abdullah F, et al. Evaluation of the Cognitive Effect of Newborn Resuscitation Training on Health-care Workers in Selected States in Northern Nigeria. *Ann Afr Med* 2018; 17: 33–9.
5. WHO. Maternal, Newborn, Child and Adolescent health. Available: [Online] 2018 [Cited 2021 Jan 15] Available from: URL: <http://www.who.int/maternal-child-adolescent/2018/en/>
6. Trevisanuto D, Bertuola F, Lanzoni P, Cavallin F, Matediana E, Manzungu OW, et al. Effect of a Neonatal Resuscitation Course on Healthcare Providers' Performances Assessed by Video Recording in a Low-Resource Setting. *PLoS ONE* 2015; 10: e0144443.
7. Mazza A, Cavallin F, Cappellari A, Divisic A, Grbin I, Akakpo J, et al. Effect of a short training on neonatal face-mask ventilation performance in a low resource setting. *PLoS ONE* 2017; 12: e0186731.
8. Duff JP, Topjian A, Berg MD, Chan M, Haskell SE, Joyner Jr BL, et al. 2018 American Heart Association Focused Update on Pediatric Advanced Life Support: An Update to the American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2018; 138: 731–9.
9. WHO. Guidelines on Basic Newborn Resuscitation. Geneva, Switzerland: 2012. ISBN-13:978-92-4150369-3.
10. Versantvoort MD, Kleinhout MY, Ockhuijsen DL, Bloemenkamp K. Helping Babies Breathe and its effects on intrapartum-related stillbirths and neonatal mortality in low-resource settings: a systematic review. *Arch Dis Child* 2020; 105: 127–33.
11. Patel A, Khatib MN, Kurhe K, Bhargava S, Bang A. Impact of neonatal resuscitation trainings on neonatal and perinatal mortality: a systematic review and meta-analysis. *BMJ Paediatr Open*. 2017; 1: e000183
12. Xu T, Wang H, Gong L, Ye H, Yu R, Wang D, et al. The impact of an intervention package promoting effective neonatal resuscitation training in rural China. *Resuscitation* 2014; 85: 253–9.
13. Hingleya S, Booth A, Hodgson J, Langworthy K, Shimizu N, Maconochie I. Concordance between the 2010 and 2015 Resuscitation Guidelines of International Liaison Committee of Resuscitation Councils (ILCOR) members and the ILCOR Consensus of Science and Treatment Recommendations (CoSTRs). *Resuscitation* 2020; 151: 111–7.

14. Azzopardi D, Strohm B, Linsell L, Hobson A, Juszczak E. Implementation and Conduct of Therapeutic Hypothermia for Perinatal Asphyxial Encephalopathy in the UK – Analysis of National Data. *PLoS ONE* 2012; 7: e38504.
 15. Hoban R, Bucher S, Neuman I, Chen M, Tesfaye N, Spector JM. 'Helping babies breathe' training in sub-saharan Africa: educational impact and learner impressions. *J Trop Pediatr* 2013; 59: 180-6.
 16. Musafili A, Essén B, Baribwira C. Evaluating Helping Babies Breathe: training for healthcare workers at hospitals in Rwanda. *Acta Pædiatr* 2013; 102: 34-8.
 17. Lai NM, Ngim CF, Fullerton PD. Teaching Medical Students Neonatal Resuscitation: Knowledge Gained and Retained from a Brief Simulation-based Training Workshop. *Educ Health (Abingdon)* 2012; 25: 105-10.
 18. Yadav S, Bhatta N, Yadav S, Kanodia P. Evaluation of Effectiveness of Neonatal Resuscitation Programme (NRP) Course among Paediatric Residents of BPKIHS. *J Nepal Paediatr Soc* 2015; 35: 18-23.
 19. Seto TL, Tabangin ME, Josyula S, Taylor KK, Vasquez JC, Kamath-Rayne BD. Educational outcomes of Helping Babies Breathe training at a community hospital in Honduras. *Perspect Med Educ* 2015; 4: 225-32.
 20. Trevisanuto D, Ferrarese P, Cavicchioli P, Fasson A, Zanardo V, Zacchello F. Knowledge gained by pediatric residents after neonatal resuscitation program courses. *Paediatr Anaesth* 2005; 15: 944-7.
 21. Bang A, Patel A, Bellad R, Gisore P, Goudar SS, Esamai F, et al. Helping Babies Breathe (HBB) training: What happens to knowledge and skills over time? *BMC Pregnancy Childbirth* 2016; 16: 364.
-