

Range for Normal Body Temperature in the General Population of Pakistan

Mehreen Adhi¹, Rabia Hasan², Fatima Noman³, Syed Faisal Mahmood⁴, Anwar Naqvi⁵, Adib-ul-Hasan Rizvi⁶
Students, Dow University of Health Sciences, Dow Medical College^{1,2}, Department of Microbiology, Liaquat National Hospital³, Department of Medicine,
The Aga Khan University Hospital⁴, Sindh Institute of Urology and Transplantation^{5,6}, Karachi, Pakistan.

Abstract

Objectives: To determine the range for normal body temperature in the general population of Pakistan and to determine if any age, sex and ambient temperature related variations exist in body temperature. Moreover, to compare how much axillary temperature differs from oral temperature measurements.

Methods: Oral as well as left and right axillary temperature recordings were made using an ordinary mercury-in-glass thermometer in 200 healthy individuals accompanying patients at various clinics at the Sindh Institute of Urology and Transplantation (SIUT) between mid-May to mid-June 2006. Data analysis was done using Epi Info version 3.3.

Results: The range for Normal Oral Temperatures fell between 97 degrees F to 99.8 degrees F (mean 98.4 degrees F). There were no significant age related ($p=0.68$) and ambient temperature related variations ($p=0.51$) in body temperature, but women had slightly higher normal temperatures than men (mean 98.5 degrees F vs. 98.3 degrees F; $p=0.01$). A wide variation existed in the difference between oral and axillary temperatures, with axillary temperatures ranging up to 2.6 degrees F lower or up to 1.1 degrees F higher than the oral temperatures (mean difference=0.85 degrees F). The correlation between oral and axillary temperatures increased at higher oral temperatures ($p=0.009$).

Conclusion: There is a range for Normal Body Temperature and any temperature above 98.6 degrees F/37 degrees C is not necessarily pathological. Women appear to have higher body temperatures. As there is no uniform oral equivalent of axillary temperature, the latter should be interpreted with caution.

Introduction

The body produces heat by muscular exercise, assimilation of food and the vital processes and this is lost from the body by radiation, conduction and vapourization, and in small amounts through urine and faeces. The balance between the heat production and heat loss determines the body temperature. Normal body function depends upon a relatively constant body temperature¹. Despite the widespread application of thermometry in clinical medicine for over a century and a half, the definition of normal body temperature is still debated².

The academic study of body temperature began in 1868³. Normal body temperature has traditionally been considered to be 98.6°F (37°C). However, a recent study indicates that normal body temperature (measured orally) varies among individuals as well as throughout the day ranging from 96°F in the morning to 99.9°F in the evening

with an overall average of 98.2°F⁴.

There are many factors causing variation in normal body temperature, for example there is a gender based variation in normal body temperature^{5,6}. Also, data is available on age related variations in the normal body temperature^{7,8}. Body temperature is very sensitive to hormone levels and women exhibit increases in body temperature of about 0.9°F at the time of ovulation⁹. In addition, exercise, digestion and underlying disorders such as chronic renal failure and shock, and neuro-psychiatric disorders such as chronic depression may alter the thermoregulatory response⁹. Ambient temperature and humidity have also been shown experimentally to affect body temperature⁹.

Temperature checking is an integral part of patient care as it influences diagnosis and subsequent patient management. The temperature measurements vary

according to the site where the temperature is recorded; oral, rectal, tympanic and axillary body temperatures⁶. Of the three sites most commonly used for clinical thermometric measurement (rectum, mouth and tympanic membrane), the oral temperature measurements have long been standard in clinical practice, largely because of accessibility, but also because oral temperature responds promptly to changes in the core temperature⁹. The temperature of the sublingual pocket is especially relevant clinically, because its main artery is a branch of the external carotid artery and, like its parent artery, responds quickly to changes in the core temperature⁹. Axillary temperature measurements are also used as an alternative being particularly convenient in young children and in uncooperative adults. However, studies have shown mixed results with regard to the oral equivalent of axillary temperatures¹⁰, and the reliability of axillary temperature measurements¹¹.

In Pakistan, the range for normal body temperature has not been ascertained and often temperature of 99°F is considered as fever. We therefore carried out a study to determine the range for normal body temperature and to determine if any age, sex and ambient temperature related variations exist in the body temperature. We was also compared how much the axillary temperatures differed from oral temperature measurements.

Subjects and Methods

We carried out a small scale observational cross sectional study in which 200 subjects including equal number of males and females were selected by stratified sampling method. All apparently healthy individuals accompanying patients at the different clinics at the Sindh Institute of Urology and Transplantation (SIUT) between mid May to mid June 2006 were included in the study. A questionnaire was prepared which contained details about the age, sex, ambient temperature, time of the day, pregnancy and last menstrual period (LMP) in case of females, history of any chronic illness like diabetes and hypertension and history of any recent infections and if the subject was on any medications, and the oral and right and left axillary temperature recordings.

Infants and young children were excluded due to non-compliance, while subjects with any chronic illness or those having a recent history of infections such as urinary tract infection (UTI), respiratory infections or sore throat etc were also excluded to avoid discrepancies in temperature recordings. Similarly, pregnant women and women suspected to be at or around their ovulation were also excluded to avoid discrepancies in temperature recordings due to altered hormone levels.

Each subject's body temperature was recorded by an ordinary mercury-in-glass thermometer. The temperature was first taken orally by positioning the bulb of the thermometer in the sublingual pocket in each subject. For axillary temperature measurements, the bulb of the thermometer was placed midway between the anterior and posterior axillary folds first in the right and then in the left axilla, and the temperature recordings for each axilla were recorded. The average of the temperatures on both sides was recorded as the mean axillary temperature in each subject for accuracy in axillary temperature measurement. For each temperature measurement the thermometer was kept at the body site for 2 minutes. The thermometer bulb was wiped with 70% alcohol at room temperature before checking the temperatures at different sites. All temperature recordings were made in the day time (morning/afternoon) in the waiting areas of the clinics at SIUT. The ambient temperature of the waiting areas was measured by using a wall mounted thermometer with a measuring range between -20°F and 140°F. This thermometer was mounted at a central point in the waiting areas each day and the ambient temperature was recorded before each subject's body temperature recording was made, so that large fluctuations in the environmental temperature do not interfere with the body temperature recordings.

Permission and approval for the study was taken from the Director of the institute prior to initiation of the study. Informed consent was taken from each subject and confidentiality and anonymity of the record was maintained. Data entry and analysis was done on Epi Info Version 3.3. The t-test was used for significance testing and a p value less than 0.05 was taken to represent a statistically significant result.

Results

A total of 200 subjects were included with equal number of males and females. Of these 16 were excluded due to history of recent infections (UTI, respiratory infections and sore throat). Of the remaining 184 subjects 52.7% (n=97) were females while 47.2% (n=87) were males. The average age of the subjects was 34 years (range=9-70 years). Using 34 years as the cutoff point, the subjects were divided into two age groups. The younger age group comprised 50% of the subjects (n=92) while the older age group also comprised 50% of the subjects (n=92). The ambient temperature of the waiting areas of the clinics ranged between 79°F and 90°F (mean=82.9 ± 1.9°F). The range for Normal Oral Temperatures for all subjects fell between 97°F to 99.8°F with the mean falling at 98.4 ± 0.52 °F. (Figure)

There were no significant age related variations in body temperature (p=0.68), but women had slightly higher




Figure: Distribution of oral temperatures in healthy individuals. Graphical representation of the distribution of oral temperatures in healthy individuals of our study population, depicting the mean oral temperature, the median and mode. The normal oral temperatures in our study population ranged from 97°F to 99.8°F.

normal body temperatures than men ($p=0.01$). The varying ambient temperature did not cause any significant variations in the body temperature ($p=0.51$). (Table-1)

Table-1: Age, Sex and Ambient Temperature related variations in normal body temperature.

		Mean Temperature (°F)	p value
Age	< 34 years	98.5	0.68
	≥ 34 years	98.4	
Sex	Male	98.3	0.01
	Female	98.5	
Ambient Temperature (°F)	< 82.5	98.5	0.51
	≥ 82.5	98.4	

The axillary temperature in the right axilla ranged from 96°F to 99.6°F (mean= $97.5 \pm 0.72^\circ\text{F}$) and that in the left axilla ranged from 96°F to 99.4°F (mean= $97.6 \pm 0.74^\circ\text{F}$). There was no significant difference in the axillary temperatures in the right and the left axilla ($p=1.00$). The average axillary temperature was therefore calculated, the mean of which was $97.5 \pm 0.68^\circ\text{F}$ ranging from 96.1°F to 99.5°F. This was lower than the oral temperature by an average of 0.85°F. However, there was a wide variation in this difference, with axillary temperatures up to 2.6°F lower or up to 1.1°F higher than the oral temperatures. About 11.4% of the subjects ($n=21$) had higher axillary temperatures while 88.5% of the subjects ($n=163$) had higher oral temperatures.

With increasing oral temperatures, the difference between the oral and axillary temperature also increased ($p=0.009$). This difference ranged from a mean of 0.5°F at oral temperatures between 97°F to 97.7°F increasing to a mean of 1.4°F at oral temperatures between 99.2°F to 99.8°F, thus showing no constant difference between the oral and axillary temperatures. (Table-2)

Table-2: Mean difference between oral and axillary temperatures with increasing oral temperatures.

	ORAL TEMPERATURES (°F)			
	97-97.7	97.8-98.4	98.5-99.1	99.2-99.8
Number of Subjects (n)	16	74	77	17
Mean Oral Temperature (°F)	97.4	98.1	98.7	99.3
	± 0.24	± 0.21	± 0.16	± 0.18
Mean Axillary Temperature (°F)	96.9	97.4	97.7	97.9
	± 0.65	± 0.62	± 0.62	± 0.73
Difference between mean oral & and mean axillary temperatures	0.5	0.7	1.0	1.4

Discussion

The main aim of this study was to determine the range for normal body temperature in apparently healthy subjects and to prove that no single cut off temperature can be ascertained to separate normal temperature from fever. This was proved by a wide variation in the normal body temperatures seen in our study. A temperature of 98.6°F is often considered as "normal" and is taken as the upper limit of normal body temperature. But one of the largest studies of oral temperature in healthy subjects demonstrated a mean temperature of 98.2°F (36.8°C) and concluded that 98.6°F (37°C) should be abandoned as a concept relevant to clinical thermometry and 98.9°F (37.2°C) in the early morning and 99.9°F (37.7°C) overall should be regarded as the upper limit of the normal oral temperature range in healthy adults aged 40 years or younger⁵. No single temperature can be considered normal, because measurements in many healthy individuals have shown a range of normal temperatures measured orally from less than 97°F (36°C) to over 99.5°F (37.5°C)¹². Our study also conflicted with the concept of 98.6°F, as the mean oral temperature of our subjects was 98.4°F with a wide range of normal oral temperatures from 97°F to 99.8°F.

Many factors can contribute to the variation of body temperature. One study pointed towards the consideration of the gender of the subject when assessing normal body temperature as it was found that the range for normal body temperature in men was 96.2°F to 100°F (35.7°C to 37.8°C) while in women it was somewhat wider, ranging from 91.7°F to 100.5°F (33.2°C to 38.1°C)⁶. Another study confirmed that women had slightly higher normal temperatures than men⁵. Wunderlich and Seguin also maintained that women have slightly higher normal temperatures than men overall and often show greater and more sudden changes in temperature⁹. This was also seen in our study as we found a statistically significant

difference between the normal temperatures of men and women, with women having slightly higher normal temperatures than men.

Some studies have been particularly conducted to determine the effect of age on the normal body temperatures. One such study used a noninvasive temporal artery thermometer to establish mean temperatures for healthy infants, children and adolescents and found a mean of 98.7°F (37.1°C) for infants 0 to 2 months, 98.4°F (36.9°C) for children 3 to 47 months, 98.2°F (36.8°C) for children 4 to 9 years and 98°F (36.7°C) for adolescents 10 to 18 years showing decreasing mean temperatures with age⁸. Another study confirmed that "older is colder" and older subjects have mean oral body temperatures lower than 98.6°F⁷. Our study conflicted with these results as there were no statistically significant age related variations in the normal body temperature in our subjects. This may have been due to the relative homogeneity in the ages of our study population.

One study also revealed the effect of moderate ambient temperature variance and proved that spuriously elevated oral temperatures may occur in warm examination areas¹³. However, as most of our subjects were in a cool and covered area of the hospital, variations in the ambient temperature were minimal and this had no significant effect on the body temperatures.

Axillary temperature is commonly used because of convenience and safety and because measurement of the oral temperature may be unhygienic. The accuracy of axillary temperature has been debated. Some studies have supported axillary temperature measurement as an acceptable alternative to rectal/oral temperature measurements¹⁴⁻¹⁶, while another study obtained a correction factor of 1°F for the correct oral equivalent of axillary temperature¹⁰. The same study also showed a wide range of difference between the right and the left axillary temperature measurements and concluded that for accurate axillary temperature, the mean of the right and left axillary temperatures should be considered¹⁰. Other studies have refuted the reliability of axillary site for temperature measurement^{17,18}. Yet another study concluded that axillary temperatures may be misleading and should be abandoned in the outpatient setting¹⁹.

Loudon showed that axillary temperature measurements with mercury thermometers vary from 2.6°F lower to 1°F higher than simultaneous oral measurements⁹. Nichols and colleagues reported that axillary temperatures exhibit differences of 0°F to 4.2°F compared with oral temperature readings in adults⁹. Falzon A, et al revealed a wide variability in the difference between oral and axillary temperature and proved that the variation between oral and

axillary temperatures increases further with increasing body temperature¹¹. Our study also showed a wide range of difference between oral and axillary temperatures and an increase in the difference with increasing oral temperatures. This may be due to the known time lag in temperature elevation between the central and the core¹¹. Thus, we were unable to find a uniform correction factor between the axillary and oral temperatures making the standard practice of adding 1°F to the axillary temperature questionable.

Unfortunately, we were unable to calibrate the thermometer between each temperature measurement which may have impacted the accuracy of our results. Another limitation of our study may have been the relatively short time the thermometer was kept at the body site.

Conclusion

There is a range for normal body temperature and any temperature above 98.6°F (37°C) is not necessarily pathological. Women appear to have higher body temperatures than men but there is no effect of age and ambient temperature level on the body temperature. As the correlation between oral and axillary temperature differs at higher and lower oral temperatures and the difference between the two increases with increasing oral temperatures, it is therefore impossible to obtain an accurate correction factor. Nevertheless, axillary temperatures are indeed a convenient mode of temperature recording in children and un-cooperative adults, but as they do not reliably reflect oral temperatures, they should therefore be interpreted with caution.

References

1. Ganong WF. Review of Medical Physiology. 21st Ed. India: McGraw Hill Co; 2003.
2. Cohen J, Powderly WG. Infectious Diseases. 2nd Ed. United States: Mosby; 2004.
3. Ng DK, Lam JC, Chow KW. Childhood fever revisited. Hong Kong Med J 2002; 8:39-43.
4. Sherwood L. Human Physiology from Cells to Systems. 5th Ed. India: Brookes/Cole-Thomson Learning; 2004.
5. Mackowiak PA, Wasserman SS, Levine MM. A critical appraisal of 98.6 degrees F, the upper limit of the normal body temperature, and other legacies of Carl Reinhold August Wunderlich. JAMA 1992; 268:1578-80.
6. Sund-Levander M, Forsberg C, Wahren LK. Normal oral, rectal, tympanic and axillary body temperature in adult men and women: a systematic literature review. Scand J Caring Sci 2002; 16:122-8.
7. Gomolin IH, Aung MM, Wolf-Klein G, Auerbach C. Older is colder: temperature range and variation in older people. J Am Geriatr Soc 2005; 53:2170-2.
8. Roy S, Powell K, Gerson LW. Temporal artery temperature measurements in healthy infants, children, and adolescents. Clin Pediatr (Phila) 2003; 42:433-7.
9. Mackowiak PA. Temperature regulation and pathogenesis of fever. In: Mandell GL, Bennett's JE, Dolin R. Douglas and Bennett's Principles and Practice of Infectious Diseases. 6th Ed. Spain: Elsevier-Churchill Livingstone; 2005, p 703-717.
10. Singh V, Sharma A, Khandelwal R, Kothari K. Variation of axillary temperature and its correlation with oral temperature. J Assoc Physicians India 2000; 48:898-900.

11. Falzon A, Grech V, Caruana B, Magro A, Attard-Montalto S. How reliable is axillary temperature measurement? *Acta Paediatr* 2003; 92:309-13.
 12. Guyton AC, Hall JE. *Textbook of Medical Physiology*. 11th Ed. India: Elsevier; 2006.
 13. Zehner WJ, Terndrup TE. The impact of moderate ambient temperature variance on the relationship between oral, rectal, and tympanic membrane temperatures. *Clin Pediatr (Phila)* 1991; 30:61-4.
 14. Lodha R, Mukerji N, Sinha N, Pandey RM, Jain Y. Is axillary temperature an appropriate surrogate for core temperature? *Indian J Pediatr* 2000; 67:571-4.
 15. Giuffre M, Heidenreich T, Carney-Gersten P, Dorsch JA, Heidenreich E. The relationship between axillary and core body temperature measurements. *Appl Nurs Res* 1990; 3:52-5.
 16. Perera KM, Colombage GR, Perera C. Axillary temperature in relation to oral temperature in a group of children in Sri Lanka. *Indian J Physiol Pharmacol* 1981; 25:26-32.
 17. Osinusi K, Njinyam MN. Comparison of body temperatures taken at different sites and the reliability of axillary temperature in screening for fever. *Afr J Med Med Sci* 1997; 26:163-6.
 18. Haddock BJ, Merrow DL, Swanson MS. The falling grace of axillary temperatures. *Pediatr Nurs* 1996; 22:121-5.
 19. Ogren JM. The inaccuracy of axillary temperatures measured with an electronic thermometer. *Am J Dis Child* 1990; 144:109-11.
-