

Breast cancer risk factors: A comparison between pre-menopausal and post-menopausal women

Zeeshan Butt,¹ Syed Furqan Haider,² Shumaila Arif,³ Muhammad Raza Khan,⁴ Umair Ashfaq,⁵
Umer Shahbaz,⁶ Mulazam Hussain Bukhari⁷

Mayo Hospital,^{1,3,5,6} Final Year MBBS, King Edward Medical University,^{2,4}
Department of Pathology, King Edward Medical University,⁷ Lahore, Pakistan.

Abstract

Objective: To evaluate the difference between pre- and post-menopausal breast cancer regarding menstrual and reproductive risk factors

Methods: The case-control study was conducted in Mayo Hospital, Lahore, between October 2008 and April 2009. Cases were breast cancer patients admitted in the surgical and oncology floors of the hospital. Controls were hospital patients without breast cancer. The study enrolled 150 cases and 300 controls. Menstrual and reproductive history was taken from both the cases and the controls. For every risk factor age-adjusted odds ratio (OR) and 95% confidence interval (CI) was calculated by logistic regression analysis, separate for pre- and post-menopausal women.

Results: Among the breast cancer patients, 42.7% were pre-menopausal and 57.3% were post-menopausal. Age at menarche had no association with breast cancer for both pre- and post-menopausal women. Nulliparity was a risk factor for both pre-menopausal (OR 4.1, 95% CI 1.4-12.6) and post-menopausal breast cancer (OR 8.2, 95% CI 1.7-40.4). Among parous women only post-menopausal females having <3 children were at increased risk for breast cancer (OR 4.5, 95% CI 1.8-11.5) compared with females having ≥ 3 children. Younger age at first live birth decreased breast cancer risk in both pre- and post-menopausal women. Breastfeeding was not associated with both pre- and post-menopausal breast cancer.

Conclusions: Majority of risk factors for pre-menopausal breast cancer are also associated with post-menopausal breast cancer except less parity, which increased the risk for post-menopausal breast cancer only.

Keywords: Breast cancer, Epidemiology, Case-control study, Risk factors (JPMA 62: 120; 2012).

Introduction

With 1 million new cases in the world each year, breast cancer is the most commonest malignancy in women and comprises 18% of all female cancers.¹ Breast cancer incidence rates are increasing in most regions of the world, especially in the developing nations.² Breast cancer is the most common female cancer in Pakistan with an age-adjusted incidence rate of 50.1 per 100,000 per year.³ Globally there is a wide variation in age standardised breast cancer incidence across different populations ranging from 19.3 per 100,000 in Eastern Africa to 89.9 per 100,000 women in Western Europe.⁴ This variation is mainly due to differences in reproductive and menstrual characteristics.

Reproductive and menstrual factors, also called hormonal factors, are the most important risk factors for breast cancer. Younger age at menarche, Nulliparity, older age at first live birth and no breastfeeding have been consistently found to increase breast cancer risk.⁵⁻²⁰

Menopause marks a crucial change in the reproductive life of a woman, bringing a decline in the levels of female sex hormones particularly oestrogen and progesterone. There is also a marked increase in the levels of gonadotrophin hormones as a result of lack of feedback inhibition by the female sex hormones. Elevated FSH and LH levels lead to stromal stimulation of the ovary, with a resultant increase in estrone levels and a decrease in estradiol levels. Inhibin levels also drop during this time because of the negative feedback of elevated FSH levels.²¹ As hormonal factors play a significant role in causing breast cancer and the onset of menopause changes hormonal environment of females, therefore pre-menopausal women might not have the same hormonal risk factors as do the post-menopausal women. Many studies have tried to address this question and have found variable results.^{5-9,11,12,14,16-19,22-23} A cohort study showed that younger age at menarche increased breast cancer risk only in pre-menopausal women, while Nulliparity increased the risk only in post-menopausal women. Older age at first live birth increased breast cancer risk in both the menopausal groups.⁷ Association of reproductive factors with breast cancer in Pakistani females, regardless of menopausal status, has been described in our previous study.²⁴ The current study presented comparison between pre-menopausal and post-menopausal women regarding breast cancer risk associated with reproductive and menstrual factors.

Patients and Methods

It was a case-control study conducted in Mayo

Hospital, Lahore, between October 2008 and April 2009. Cases were breast cancer patients admitted in the surgical and oncology floors of the hospital. Controls were hospital patients without breast cancer. Cancer patients were not eligible as controls. As less cases than controls were available, it was decided to take cases and controls in a ratio of 1:2. Assuming 5% of control population to be nulliparous, 141 cases and 282 controls were needed to detect the odds ratio of 3 with power of 80% and confidence limits of 95%. We enrolled 150 cases and 300 controls. Cases were enrolled through consecutive convenience sampling, and controls through quota sampling according to pre-specified age groups.

After obtaining informed consent, both the cases and the controls were asked about their basic demographic factors. Menstrual and reproductive history was taken, including age at menarche, menopausal status, age at menopause, number of children born alive, age at first live birth and breastfeeding practice. Menopause was defined as absence of menstruation for at least six months before the diagnosis of breast cancer for cases and date of admission for controls.

Logistic regression was used to calculate age-adjusted odds ratio and 95% confidence interval for every hormonal factor, separate for pre- and post-menopausal women. The p-value was calculated using Chi-square test. All analyses were done with SPSS version 16. The study design was approved by the Ethical Review Board of the King Edward Medical University, Lahore.

Results

Among the breast cancer patients 42.7% were pre-menopausal and 57.3% were post-menopausal. Age at menarche had no association with breast cancer for both pre- and post-menopausal women. Nulliparity was a risk factor for both pre- and post-menopausal breast cancer, but the association was stronger for post-menopausal women. Higher age at menopause did not increase breast cancer risk in post-menopausal women (Table-1).

Among parous post-menopausal women, breast cancer OR was 4.5 (95% CI 1.8-11.5) for women having <3 children compared with women having ≥ 3 children. No association was found between less parity and pre-menopausal breast cancer (Table-2). Among pre-menopausal parous women more cases than controls (13.0% vs. 0%) had age ≥ 30 years at first live birth (p <0.001). History of breastfeeding did not increase breast cancer risk for both pre- and post-

Table-1. Breast cancer risk factors for all women.

Variables	Case N (%)	Control N (%)	OR*	95% CI	P-value
Pre-Menopausal women					
Age at Menarche					
<12	6 (9.4)	4 (3.6)	2.7	0.7-10.2	0-13
≥ 12	58 (90.6)	107 (96.4)			
Parity					
Nulliparous	10 (15.6)	6 (5.4)	4.1	1.4-12.6	0.01
Parous	54 (84.4)	105 (94.6)			
Post-Menopausal women					
Age at menarche					
<12	1 (1.2)	5 (2.6)	0.4	0.04-3.7	0.43
≥ 12	85 (98.8)	184 (97.4)			
Parity					
Nulliparous	7 (8.1)	2 (1.1)	8.2	1.7-40.4	0.01
Parous	79 (91.9)	187 (98.9)			
Age at Menopause					
≥ 50	17 (19.8)	36 (19.0)	1.0	0.5-2.0	0.97
< 50	69(80.2)	153 (81.0)			

* Age-Adjusted Odds Ratio (OR).

Table-2: Breast cancer risk factors for parous women.

Variables	Case N (%)	Control N (%)	OR*	95% CI	P-value
Pre-Menopausal women					
No of Live Births					
< 3	8(14.8)	14(13.3)	1.6	0.6-4.5	0.35
≥ 3	46(85.2)	91(86.7)			
Age at First Live Birth					
< 30	47 (87.0)	105 (100)	†		< 0.01
≥ 30	7 (13.0)	0 (0.0)			
Lactation					
Never	4 (7.4)	6 (5.7)	1.4	0.4-5.3	0.61
Ever	50(92.6)	99(94.3)			
Post Menopausal women					
No of live Births					
< 3	13 (16.5)	8 (4.3)	4.5	1.8-11.5	< 0.01
≥ 3	66 (83.5)	179 (95.7)			
Age at First Live Birth					
< 30	71 (89.9)	180 (96.3)	0.3	0.12-0.99	0.048
≥ 30	8 (10.1)	7 (3.7)			
Lactation					
Never	3 (3.8)	3 (1.6)	2.5	0.5-12.6	0.27
Ever	76 (96.2)	184 (98.4)			

* = Age-Adjusted Odds Ratio (OR).

† = OR could not be calculated. P-value was calculated using chi-square test.

menopausal women.

Discussion

The study found that younger age at menarche did not increase breast cancer risk for both pre- and post-menopausal women while Nulliparity was a risk factor for both menopausal groups. Higher age at menopause was not a risk factor for post-menopausal women.

Previous studies have reported inconsistent results regarding age at menarche and breast cancer risk. Lubin et

al⁸ found that younger age at menarche increased breast cancer risk only in pre-menopausal women, while Tung et al²³ reported increased risk only for post-menopausal women. In some studies, age at menarche was associated with both pre- and post-menopausal breast cancer^{11,16} while in another study it had no association with either pre- or post-menopausal breast cancer.¹² Majority of these studies relied on patient's recall for determining age at menarche. Correct recall of events that happened years ago depends on many factors, including age, education, and mental well-

being. Recall-bias could be the reason for these inconsistent findings about age at menarche and breast cancer risk. A review article showed that age at menarche was associated with both pre- and post-menopausal breast cancer, but the association was stronger for pre-menopausal women. For each additional year of age at menarche, the risk of female breast cancer decreased by 9% for pre-menopausal women, and by 4% for post-menopausal women.⁶ Similar findings were found in a study which showed that younger age at menarche explained 70% of pre-menopausal breast cancers while it explained only 27.5% of post-menopausal breast cancers.¹⁶

We found that both pre- and post-menopausal nulliparous women were at increased risk for developing breast cancer. A previous study has found that Nulliparity is associated with only post-menopausal breast cancer.⁵ However, another study has reported increased risk for only pre-menopausal women.²² A review article showed that pre-menopausal breast cancer risk was reduced by 3% per full-term pregnancy while the risk reduction was 12% for post-menopausal breast cancer.⁶ It seems that the protective effect of parity against breast cancer is more for post-menopausal women than for pre-menopausal women.

In a previous study, it was found that women who gave birth to a child before 24 years of age exhibit a decrease in their lifetime risk of developing breast cancer, and moreover additional pregnancies increase the protective effect. The breast tissue of normally cycling women contains three identifiable types of lobules; the undifferentiated Lobules type 1 (Lob 1) and the more developed Lobules type 2 and Lobules type 3. The breast attains its maximum development during pregnancy and lactation (Lobules type 4). After menopause the breast regresses in both nulliparous and parous women and consists only of Lobules type 1. Despite the similarity in the lobular composition of the breast at menopause, in both the parous and nulliparous women, Lob 1 in these two groups of women are different biologically, and exhibit different susceptibility to carcinogenesis. It was suggested that Lob 1 found in the breast of nulliparous women and of parous women with breast cancer never went through the process of differentiation, retaining a high concentration of epithelial cells that are targets for carcinogens and are, therefore, susceptible to undergo neoplastic transformation. These epithelial cells are called Stem cells 1, whereas Lob 1 structures found in the breast of early parous post-menopausal women free of mammary pathology, on the contrary, are composed of an epithelial cell population that is refractory to transformation, called Stem cells 2.¹³

In our study, increasing parity decreased breast

cancer risk only in post-menopausal women while younger age at first live birth (FLB) decreased risk in both pre- and post-menopausal women. The findings about decreased risk for increasing parity are consistent with some previous studies.¹⁴⁻¹⁶ Some studies have found that increasing parity decreased risk in both pre- and post-menopausal women.^{19,22} In a local study increasing parity reduced breast cancer risk only in pre-menopausal women.¹⁷

Our findings about age at FLB are consistent with many previous studies which found that younger age at FLB decreased breast cancer risk in both pre- and post-menopausal women.^{6,7,16,22} However, some studies found that a higher age at first live birth is protective only for postmenopausal women.^{8,9,19} In a cohort study relative risk of breast cancer for age at FLB >30 years was 1.63 for pre-menopausal women and 1.35 for post-menopausal women.⁷

Younger age at FLB reduces breast cancer risk in two ways. First, early pregnancy results in early differentiation of mammary gland cells. Second, women with early age at FLB usually have longer cumulative duration of breastfeeding which further reduces breast cancer risk. Some studies have found that younger age at FLB is more important than increasing parity in reducing breast cancer risk. Gao showed that increasing parity no longer decreased breast cancer risk after adjusting for age at FLB.¹⁶

We found that breastfeeding was not protective against breast cancer for both pre- and post-menopausal women. Our result is consistent with a local study,¹⁷ but majority of Western studies have shown that breastfeeding is protective against pre-menopausal breast cancer.^{18,19} However, some studies found protective effect only for post-menopausal women.¹⁶ A collaborative re-analysis of 47 studies showed that breastfeeding was protective for both premenopausal and post menopausal women.²⁵

Our study had some limitations. First, we studied prevalent breast cancer cases rather than incident ones. Second, the sample size was small which resulted in wider confidence intervals. We found only one local study similar to ours that assessed role of reproductive factors in causing breast cancer separate for pre- and post-menopausal women.

Conclusion

Despite its limitations, the study will be a useful addition to current understanding of epidemiology of breast cancer in Pakistan. Studies done on a wider scale are needed to establish the findings of the study related to risk factors for pre-menopausal and post-menopausal breast cancer in the country.

References

1. McPherson K, Steel CM, Dixon JM. ABC of breast diseases. Breast cancer-epidemiology, risk factors, and genetics. *BMJ* 2000; 321: 624-8.
2. Bray F, McCarron P, Parkin DM. The changing global patterns of female breast cancer incidence and mortality. *Breast Cancer Res* 2004; 6: 229-39.
3. Sohail S, Alam SN. Breast cancer in Pakistan - awareness and early detection. *J Coll Physicians Surg Pak* 2007; 17: 711-2.
4. Ferlay J, Shin H, Bray F, Forman D, Mathers C, Parkin D. GLOBOCAN 2008. Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10 [Internet]. Lyon: International Agency for Research on Cancer 2010.
5. Lee HP, Gourley L, Duffy SW, Esteve J, Lee J, Day NE. Risk factors for breast cancer by age and menopausal status: a case-control study in Singapore. *Cancer Causes Control* 1992; 3: 313-22.
6. Clavel-Chapelon F, Gerber M. Reproductive factors and breast cancer risk. Do they differ according to age at diagnosis? *Breast Cancer Res Treat* 2002; 72: 107-15.
7. Clavel-Chapelon F; EBM-EPIC Group. Differential effects of reproductive factors on the risk of pre- and postmenopausal breast cancer. Results from a large cohort of French women. *Br J Cancer* 2002; 86: 723-7.
8. Lubin JH, Burns PE, Blot WJ, Lees AW, May C, Morris LE, et al. Risk factors for breast cancer in women in northern Alberta, Canada, as related to age at diagnosis. *J Natl Cancer Inst* 1982; 68: 211-7.
9. Stavrakys K, Emmons S. Breast cancer in premenopausal and postmenopausal women. *J Natl Cancer Inst* 1974; 53: 647-54.
10. Döntenwill W. [Significance of follicular hormone on the pathogenesis of carcinoma of the breast and the relation of hormone-induced breast disease to cancer of the breast; statistical research on the problem of the development of breast cancer from breast disease.]. *Z Krebsforsch* 1955; 60: 476-81.
11. Hsieh CC, Trichopoulos D, Katsouyanni K, Yuasa S. Age at menarche, age at menopause, height and obesity as risk factors for breast cancer: associations and interactions in an international case-control study. *Int J Cancer* 1990; 46: 796-800.
12. Beiler JS, Zhu K, Hunter S, Payne-Wilks K, Roland CL, Chinchilli VM. A case-control study of menstrual factors in relation to breast cancer risk in African-American women. *J Natl Med Assoc* 2003; 95: 930-8.
13. Russo J, Moral R, Balogh GA, Mailo D, Russo IH. The protective role of pregnancy in breast cancer. *Breast Cancer Res* 2005; 7: 131-42.
14. Antoniou AC, Shenton A, Maher ER, Watson E, Woodward E, Lalloo F, et al. Parity and breast cancer risk among BRCA1 and BRCA2 mutation carriers. *Breast Cancer Res* 2006; 8: R72.
15. Lee E, Ma H, McKean-Cowdin R, Van Den Berg D, Bernstein L, Henderson BE, Ursin G. Effect of reproductive factors and oral contraceptives on breast cancer risk in BRCA1/2 mutation carriers and noncarriers: results from a population-based study. *Cancer Epidemiol Biomarkers Prev* 2008; 17: 3170-8.
16. Gao YT, Shu XO, Dai Q, Potter JD, Brinton LA, Wen W, et al. Association of menstrual and reproductive factors with breast cancer risk: results from the Shanghai Breast Cancer Study. *Int J Cancer* 2000; 87: 295-300.
17. Gilani GM, Kamal S. Risk factors for breast cancer in Pakistani women aged less than 45 years. *Ann Hum Biol* 2004; 31: 398-407.
18. Newcomb PA, Storer BE, Longnecker MP, Mittendorf R, Greenberg ER, Clapp RW, et al. Lactation and a reduced risk of premenopausal breast cancer. *N Engl J Med* 1994; 330: 81-7.
19. Ursin Gea, Bernstein L, Wang Y, Lord SJ, Deapen D, Liff JM, et al. Reproductive factors and risk of breast carcinoma in a study of white and African-American women. *Cancer* 2004; 101: 353-62.
20. Enger SM, Ross RK, Henderson B, Bernstein L. Breastfeeding history, pregnancy experience and risk of breast cancer. *Br J Cancer* 1997; 76: 118-23.
21. Lenton E, de Kretser D, Woodward A, Robertson D. Inhibin concentrations throughout the menstrual cycles of normal, infertile, and older women compared with those during spontaneous conception cycles. *J Clin Endocrinol Metab* 1991; 73: 1180-90.
22. Gajalakshmi CK, Shanta V. Risk factors for female breast cancer. A hospital-based case-control study in Madras, India. *Acta Oncol* 1991; 30: 569-74.
23. Tung HT, Tsukuma H, Tanaka H, Kinoshita N, Koyama Y, Ajiki W, et al. Risk factors for breast cancer in Japan, with special attention to anthropometric measurements and reproductive history. *Jpn J Clin Oncol* 1999; 29: 137-46.
24. Butt Z, Shahbaz U, Hashmi AT, Naseem T, Khan MZ, et al. Frequency of conventional risk factors in patients with acute coronary syndrome in males and females. *Ann King Edward Med Uni* 2010; 16: 55-8.
25. Collaboration Group on Hormonal Factors in Breast Cancer. Breast cancer and breastfeeding: collaborative re-analysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet* 2002; 360: 187-95.